



GENVOYA® MADE SIMPLE

ONE COMPLETE HIV REGIMEN - TAKE ONCE DAILY WITH FOOD

1

TAKE IT WITH FOOD

Take one tablet once daily with food. Food is required for reliable absorption.

2

DO NOT STOP ON YOUR OWN

NO UNPLANNED GAPS

Stopping can allow HIV to rebound. If you have hepatitis B, stopping may cause a serious liver flare.

3

ONE COMPLETE REGIMEN

Genvoya contains four medicines. Do not add or remove HIV medicines unless your specialist directs it.

4

CHECK EVERY NEW MEDICINE

Cobicistat causes many interactions. The pharmacist must check every prescription, OTC medicine, and supplement.

5

SEPARATE ANTACIDS

Antacids containing aluminum or magnesium should generally be separated from Genvoya. Ask for exact timing.

6

KEEP LABS & REFILLS ON TIME

Kidney function, liver health, viral load, and hepatitis B status require regular follow-up.

MY DAILY GENVOYA PLAN

My daily meal/time: _____

My prescriber: _____

My next refill date: _____

MISSED A DOSE?

Take it when you remember unless it is nearly time for the next dose. Do not double. Call if you are unsure or may run out.

CALL RIGHT AWAY FOR

- Yellow skin or eyes, dark urine, or severe abdominal pain
- Very little urine, swelling, or severe weakness
- Rash with face, lip, tongue, or throat swelling
- Severe muscle pain or unusual weakness
- Any risk of running out or stopping

FOOD, ANTACIDS & INTERACTIONS

TAKE WITH FOOD

A snack or meal is required. Build the dose into a reliable daily eating routine.

ANTACIDS & MINERALS

Ask Fairview about aluminum, magnesium, calcium, or iron products and exact timing.

HIGH-INTERACTION REGIMEN

Never start rifampin, seizure medicines, St. John's wort, cholesterol medicines, or other prescriptions without a check.

IMPORTANT SAFETY & SUCCESS REMINDERS

HEPATITIS B

Testing matters before treatment. Stopping may cause a serious hepatitis B flare.

KIDNEY MONITORING

Keep kidney labs and report reduced urine, swelling, or unusual weakness.

COMPLETE REGIMEN

Do not combine with other HIV medicines unless your HIV specialist directs it.

PRIVACY & SUPPORT

Fairview helps protect privacy, prevent refill gaps, and solve insurance barriers.



MY GENVOYA® GUIDE

KEEP THIS PAGE WITH YOUR MEDICATIONS

MY TREATMENT PLAN

My name: _____

My daily meal/time: _____

My HIV clinician: _____

Next refill date: _____

Next lab date: _____

ONE TABLET DAILY WITH FOOD - NO GAPS

IF I MISS A DOSE

- 1 Take it as soon as you remember.
- 2 If the next dose is close, skip and resume the regular schedule.
- 3 Do not take two tablets together.
- 4 Call before the last tablet if a refill gap is possible.

CHECK BEFORE TAKING

ANTACIDS

Aluminum or magnesium products may need separation.

MINERALS

Calcium or iron timing should be confirmed.

NEW MEDICINES

Cobicistat causes many important interactions.

OTHER HIV DRUGS

Genvoya is already a complete regimen.

CALL MY CARE TEAM FOR

- Yellow skin or eyes or dark urine
- Very little urine or new swelling
- Severe or persistent nausea/vomiting
- A rash or facial swelling
- A new prescription or supplement
- Any chance of missing refills

DO NOT STOP ON MY OWN

Stopping can allow HIV to rebound and may cause resistance. If hepatitis B is present, stopping can trigger a serious liver flare.

CALL BEFORE THE LAST TABLET

FAIRVIEW SUPPORT

PRIVATE COUNSELING

Questions answered discreetly.

INTERACTION CHECKS

Every prescription and supplement reviewed.

REFILL PROTECTION

Call early if travel, insurance, or supply may cause a gap.

COST SUPPORT

Coverage and assistance options checked for your prescription.