



INGREZZA MADE SIMPLE

A patient guide to once-daily valbenazine

Take it once daily and report sleepiness, new stiffness or slowness, fainting, or worsening depression promptly.

1

ONCE DAILY

Take at the same time, with or without food.

2

KNOW THE GOAL

Used for tardive dyskinesia and certain Huntington chorea symptoms.

3

WATCH SLEEPINESS

Use caution with driving and alcohol.

4

WATCH PARKINSONISM

Report stiffness, slowness, tremor, balance trouble, or falls.

5

CHECK INTERACTIONS

Some CYP3A4 or CYP2D6 medicines and other VMAT2 drugs need review.

6

URGENT SIGNS

Fainting, severe heartbeat symptoms, high fever with rigidity, or suicidal crisis needs urgent help.

YOUR DAILY INGREZZA PLAN

- My dose: _____
- My daily time: _____
- I take it for: TD / Huntington chorea
- My next follow-up: _____
- Do not double a missed dose. Follow label instructions.

IMPORTANT SAFETY REMINDERS

- Huntington disease patients and families should monitor depression and suicidal thinking.
- Tell the team about heart-rhythm problems, liver disease, pregnancy, or breastfeeding.
- Do not combine with other VMAT2 inhibitors unless directed.
- Movement improvement may take time; keep follow-up appointments.

HOW FAIRVIEW HELPS

- Appointment and refill reminders
- Medication interaction review
- Side-effect and movement screening
- Cost and coverage support
- Private, nonjudgmental counseling



FAIRVIEW PHARMACY

INGREZZA - QUICK GUIDE

Simplified refrigerator reference

MY MEDICATION PLAN

My name: _____

My dose: _____

My schedule / interval: _____

My next appointment or refill: _____

My prescriber / clinic: _____

THE ONE RULE TO REMEMBER

Take it once daily and report sleepiness, new stiffness or slowness, fainting, or worsening depression promptly.

CALL THE CARE TEAM PROMPTLY FOR

- Report stiffness, slowness, tremor, balance trouble, or falls.
- Some CYP3A4 or CYP2D6 medicines and other VMAT2 drugs need review.
- Fainting, severe heartbeat symptoms, high fever with rigidity, or suicidal crisis needs urgent help.
- Huntington disease patients and families should monitor depression and suicidal thinking.
- Tell the team about heart-rhythm problems, liver disease, pregnancy, or breastfeeding.

FAIRVIEW PHARMACY 601-544-4871

Call before a missed injection becomes overdue or before changing a daily dose.

Clinically reviewed: July 2026

This guide does not replace instructions from your prescriber or clinic.