



INVEGA SUSTENNA MADE SIMPLE

A patient guide to monthly paliperidone injections

Keep every monthly injection appointment. If a dose is late, call the clinic - the restart plan depends on how long it has been.

1

KEEP THE DATE

Usually given once monthly by a healthcare professional.

2

KNOW YOUR START

Early loading injections follow a special schedule. Do not miss them.

3

CALL IF LATE

Never guess or restart on your own after a missed injection.

4

WATCH MOVEMENT

Report stiffness, tremor, restlessness, or new uncontrolled movements.

5

CHECK BODY HEALTH

Weight, glucose, cholesterol, blood pressure, and prolactin effects may need monitoring.

6

GET URGENT HELP

High fever with rigid muscles, confusion, fainting, or severe allergic symptoms need urgent care.

YOUR MONTHLY INJECTION PLAN

- My dose: _____
- My injection date each month: _____
- My next appointment: _____
- Clinic phone: _____
- If I miss an appointment: call the clinic the same day.

IMPORTANT SAFETY REMINDERS

- Do not use for dementia-related psychosis unless specifically prescribed for an approved reason.
- Tell the team about pregnancy, kidney problems, fainting, seizures, or swallowing difficulty.
- Rise slowly; dizziness and falls can occur.
- Do not drive until you know how the injection affects you.

HOW FAIRVIEW HELPS

- Appointment and refill reminders
- Medication interaction review
- Side-effect and movement screening
- Cost and coverage support
- Private, nonjudgmental counseling



FAIRVIEW PHARMACY

INVEGA SUSTENNA - QUICK GUIDE

Simplified refrigerator reference

MY MEDICATION PLAN

My name: _____

My dose: _____

My schedule / interval: _____

My next appointment or refill: _____

My prescriber / clinic: _____

THE ONE RULE TO REMEMBER

Keep every monthly injection appointment. If a dose is late, call the clinic - the restart plan depends on how long it has been.

CALL THE CARE TEAM PROMPTLY FOR

- Report stiffness, tremor, restlessness, or new uncontrolled movements.
- Weight, glucose, cholesterol, blood pressure, and prolactin effects may need monitoring.
- High fever with rigid muscles, confusion, fainting, or severe allergic symptoms need urgent care.
- Do not use for dementia-related psychosis unless specifically prescribed for an approved reason.
- Tell the team about pregnancy, kidney problems, fainting, seizures, or swallowing difficulty.

FAIRVIEW PHARMACY 601-544-4871

Call before a missed injection becomes overdue or before changing a daily dose.

Clinically reviewed: July 2026

This guide does not replace instructions from your prescriber or clinic.