



INVEGA TRINZA MADE SIMPLE

A patient guide to 3-month paliperidone injections

This injection is given every 3 months only after stable treatment with monthly Invega Sustenna. Keep the appointment window.

1

EVERY 3 MONTHS

Your clinic schedules each dose. Put the date in two places.

2

STABLE FIRST

Trinza is used only after adequate monthly Sustenna treatment.

3

CALL IF LATE

A late injection may require a different restart sequence.

4

WATCH MOVEMENT

Report stiffness, tremor, restlessness, or new uncontrolled movements.

5

MONITOR HEALTH

Weight, glucose, cholesterol, blood pressure, kidney function, and prolactin effects may be checked.

6

URGENT SIGNS

High fever with rigid muscles, confusion, fainting, or severe allergy needs urgent care.

YOUR 3-MONTH INJECTION PLAN

- My dose: _____
- My scheduled date: _____
- My allowable appointment window: ask clinic
- My next appointment: _____
- If the date may be missed, call before the appointment.

IMPORTANT SAFETY REMINDERS

- Do not switch between Sustenna, Trinza, or Hafyera without the prescriber.
- Medication remains in the body for months, so side effects may take time to improve.
- Rise slowly and use fall precautions if dizzy.
- Tell the team about kidney problems, seizures, pregnancy, or swallowing difficulty.

HOW FAIRVIEW HELPS

- Appointment and refill reminders
- Medication interaction review
- Side-effect and movement screening
- Cost and coverage support
- Private, nonjudgmental counseling



FAIRVIEW PHARMACY

INVEGA TRINZA - QUICK GUIDE

Simplified refrigerator reference

MY MEDICATION PLAN

My name: _____

My dose: _____

My schedule / interval: _____

My next appointment or refill: _____

My prescriber / clinic: _____

THE ONE RULE TO REMEMBER

This injection is given every 3 months only after stable treatment with monthly Invega Sustenna. Keep the appointment window.

CALL THE CARE TEAM PROMPTLY FOR

- Report stiffness, tremor, restlessness, or new uncontrolled movements.
- Weight, glucose, cholesterol, blood pressure, kidney function, and prolactin effects may be checked.
- High fever with rigid muscles, confusion, fainting, or severe allergy needs urgent care.
- Do not switch between Sustenna, Trinza, or Hafyera without the prescriber.
- Medication remains in the body for months, so side effects may take time to improve.

FAIRVIEW PHARMACY 601-544-4871

Call before a missed injection becomes overdue or before changing a daily dose.

Clinically reviewed: July 2026

This guide does not replace instructions from your prescriber or clinic.