



JULUCA® MADE SIMPLE

A TWO-DRUG SWITCH REGIMEN - TAKE ONCE DAILY WITH A MEAL

1

TAKE WITH A MEAL

FOOD IS REQUIRED

Take one tablet once daily with a real meal. A protein drink alone is not enough.

2

USE ONLY AS A SWITCH REGIMEN

Juluca is for selected adults already virally suppressed with no treatment failure or relevant resistance.

3

NO PROTON PUMP INHIBITORS

Omeprazole, pantoprazole, esomeprazole, and similar medicines must not be used with Juluca.

4

TIME OTHER ACID REDUCERS

H2 blockers and antacids require specific spacing. Ask Fairview for a written schedule.

5

CHECK MINERALS & NEW DRUGS

Calcium, iron, magnesium, aluminum, rifampin, seizure medicines, and St. John's wort require review.

6

KEEP LABS & REFILLS ON TIME

Viral load monitoring confirms that the simplified regimen continues to work.

MY DAILY JULUCA PLAN

My daily meal/time: _____

My HIV clinician: _____

My next refill date: _____

MISSED A DOSE?

Take it with a meal when you remember unless the next dose is close. Do not double. Never take a catch-up dose without food.

CALL RIGHT AWAY FOR

- Rash with fever, facial swelling, or trouble breathing
- Severe depression or thoughts of self-harm
- Yellow skin or eyes, dark urine, or severe abdominal pain
- New fainting, severe palpitations, or chest symptoms
- Any risk of a refill gap

ACID REDUCERS & FOOD - CRITICAL RULES

PROTON PUMP INHIBITORS

Do not use omeprazole, pantoprazole, esomeprazole, or similar PPIs with Juluca.

H2 BLOCKERS

Medicines such as famotidine require exact spacing before or after Juluca.

ANTACIDS / MINERALS

Aluminum, magnesium, calcium, or iron products also require a pharmacist timing plan.

IMPORTANT SAFETY & SUCCESS REMINDERS

MEAL REQUIRED

Take with enough food for absorption; a nutritional drink alone is not sufficient.

SWITCH REGIMEN

Do not use Juluca as initial therapy or after resistance without specialist confirmation.

MOOD MONITORING

Report severe depression, unusual mood changes, or self-harm thoughts.

PRIVACY & SUPPORT

Fairview provides discreet counseling and a written interaction schedule.



MY JULUCA® GUIDE

KEEP THIS PAGE WITH YOUR MEDICATIONS

MY TREATMENT PLAN

My name: _____

My meal/time: _____

My HIV clinician: _____

Next refill date: _____

Next viral load: _____
ONE TABLET DAILY WITH A MEAL

IF I MISS A DOSE

- 1 Take it with a meal when remembered.
- 2 If the next dose is close, skip the missed dose.
- 3 Do not double and do not take without food.
- 4 Call early if a refill gap is possible.

ACID MEDICINE RULES

PPIS

Omeprazole and similar medicines are not allowed.

H2 BLOCKERS

Famotidine and similar drugs need exact spacing.

ANTACIDS

Aluminum or magnesium products need spacing.

MINERALS

Calcium and iron timing must be checked.

CALL MY CARE TEAM FOR

- Rash, fever, or facial swelling
- Severe depression or self-harm thoughts
- Yellow skin or eyes or dark urine
- Fainting or severe palpitations
- A new acid-reducing medicine
- Any chance of missing refills

FOOD IS PART OF THE DOSE

Juluca must be taken with a meal so rilpivirine is absorbed properly. A protein drink or light beverage alone is not enough.

BUILD IT INTO A RELIABLE DAILY MEAL

FAIRVIEW SUPPORT

PRIVATE COUNSELING

Questions answered discreetly.

INTERACTION CHECKS

Every prescription and supplement reviewed.

REFILL PROTECTION

Call early if travel, insurance, or supply may cause a gap.

COST SUPPORT

Coverage and assistance options checked for your prescription.