



FAIRVIEW PHARMACY

Rooted In Care. Built For Everyone.

MYRBETRIQ® MADE SIMPLE

A patient guide to taking mirabegron safely

1

TAKE IT ONCE DAILY

Take it at the same time each day. Do not change the dose yourself.

2

SWALLOW IT WHOLE

Do not crush, chew, or split the extended-release tablet.

3

CHECK BLOOD PRESSURE

Myrbetriq can raise blood pressure. Keep your monitoring appointments.

4

WATCH URINE FLOW

Call promptly if you cannot urinate or your bladder feels painfully full.

5

CHECK NEW MEDICINES

Myrbetriq can affect some medicines, including certain heart and mental-health drugs.

6

KNOW WHEN TO GET HELP

Face or throat swelling, trouble breathing, or severe urinary retention needs urgent help.

YOUR DAILY MYRBETRIQ PLAN

My strength: _____ Daily time: _____

Why I take it: _____

What improvement may look like

Fewer sudden urges, fewer bathroom trips, and fewer leakage episodes. Improvement may build gradually over several weeks.

Do not stop early just because improvement is gradual. Call Fairview if you are unsure.

GET HELP RIGHT AWAY FOR

- Swelling of the face, lips, tongue, or throat
- Trouble breathing or swallowing
- Inability to urinate with pain or bladder fullness
- Severe chest pounding, fainting, or neurologic symptoms

CALL YOUR DOCTOR OR PHARMACIST

- Blood pressure is repeatedly high
- Fast heartbeat, palpitations, or strong headache
- Burning urination, fever, or back pain
- New constipation, dizziness, or bothersome side effects

IMPORTANT MEDICINE CHECKS

Tell Fairview if you take digoxin, metoprolol, flecainide, propafenone, certain antidepressants or antipsychotics, or another bladder medicine. Myrbetriq can increase exposure to some CYP2D6 medicines, and digoxin may need closer monitoring.

Never start, stop, or change a prescription without checking first.

FAIRVIEW PHARMACY 601-544-4871

500 Katie Avenue, Hattiesburg, MS 39401

Clinically reviewed: July 2026

General education only; follow your prescriber and pharmacist instructions.

MY MYRBETRIQ® GUIDE

KEEP THIS WITH YOUR MEDICINES

My name: _____

My strength: _____

My daily time: _____

Prescriber: _____

My blood pressure goal: _____

THE 4 RULES I SHOULD REMEMBER

- 1 Take once daily at the same time.
- 2 Swallow the tablet whole.
- 3 Check blood pressure as directed.
- 4 Call promptly if I cannot urinate.

IF I MISS A DOSE

General rule: Take the missed dose when you remember unless it is close to your next scheduled dose. Skip the missed dose if it is nearly time for the next one. Do not take two doses at the same time. Follow the instructions on your prescription label or call Fairview.

Remember: symptom improvement can take several weeks. Keep taking it unless your clinician tells you to stop.

CALL FAIRVIEW OR YOUR DOCTOR FOR

- Repeatedly high blood-pressure readings
- Fast heartbeat or new palpitations
- Burning urination, fever, or back/side pain
- Trouble starting urine or a weak stream
- Side effects that interfere with daily life

GET URGENT HELP FOR

- Face, lip, tongue, or throat swelling
- Trouble breathing or swallowing
- Painful inability to urinate
- Fainting, severe chest symptoms, or stroke-like signs

BEFORE STARTING ANY NEW MEDICINE

Tell the pharmacist that you take Myrbetriq. Important checks may include digoxin, metoprolol, flecainide, propafenone, certain antidepressants or antipsychotics, and other medicines for overactive bladder.

Questions? Fairview Pharmacy • 601-544-4871