F+ FAIRVIEW PHARMACY

REXULTI MADE SIMPLE

A patient guide to taking brexpiprazole safely

Take it once daily and know the reason it was prescribed: schizophrenia, add-on depression treatment, or agitation associated with Alzheimer dementia.

1 ONCE DAILY

Take at the same time, with or without food.

2 KNOW THE PURPOSE

Dose and monitoring differ by condition.

3 WATCH RESTLESSNESS

Report pacing, inability to sit still, stiffness, tremor, or uncontrolled movements.

4 TRACK WEIGHT & LABS

Weight, glucose, cholesterol, and blood pressure may change.

5 WATCH MOOD & IMPULSES

Report suicidal thoughts or new hard-to-control urges.

6 URGENT SIGNS

High fever, rigid muscles, confusion, fainting, stroke signs, or severe allergy needs urgent care.

YOUR DAILY REXULTI PLAN

- My dose: _____
- My daily time: _____
- Why I take it: _____
- My next follow-up: _____
- Do not increase or stop it without instructions.

IMPORTANT SAFETY REMINDERS

- Older adults with dementia-related psychosis have increased mortality risk; use only for an approved, clinician-directed reason.
- For Alzheimer-related agitation, continue non-drug support and reassess benefit.
- Use caution with driving and falls.
- Tell the pharmacist about CYP2D6 and CYP3A4 interacting medicines.

HOW FAIRVIEW HELPS

- Appointment and refill reminders
- Medication interaction review
- Side-effect and movement screening
- Cost and coverage support
- Private, nonjudgmental counseling

FAIRVIEW PHARMACY | 601-544-4871

Clinically reviewed: July 2026 | This guide supports - but does not replace - individualized care.

F+ **FAIRVIEW PHARMACY**

REXULTI - QUICK GUIDE

Simplified refrigerator reference

MY MEDICATION PLAN

My name: _____

My dose: _____

My schedule / interval: _____

My next appointment or refill: _____

My prescriber / clinic: _____

THE ONE RULE TO REMEMBER

Take it once daily and know the reason it was prescribed: schizophrenia, add-on depression treatment, or agitation associated with Alzheimer dementia.

CALL THE CARE TEAM PROMPTLY FOR

- Weight, glucose, cholesterol, and blood pressure may change.
- Report suicidal thoughts or new hard-to-control urges.
- High fever, rigid muscles, confusion, fainting, stroke signs, or severe allergy needs urgent care.
- Older adults with dementia-related psychosis have increased mortality risk; use only for an approved, clinician-directed reason.
- For Alzheimer-related agitation, continue non-drug support and reassess benefit.

FAIRVIEW PHARMACY 601-544-4871

Call before a missed injection becomes overdue or before changing a daily dose.

Clinically reviewed: July 2026

This guide does not replace instructions from your prescriber or clinic.