



TIVICAY® MADE SIMPLE

DOLUTEGRAVIR - MUST BE USED WITH THE REST OF YOUR HIV REGIMEN

1

TAKE IT AS PRESCRIBED

Many patients take it once daily, but some require twice-daily dosing. Follow the exact label.

2

NOT A COMPLETE REGIMEN

KNOW THE REST OF YOUR REGIMEN

Tivicay must be combined with other HIV medicines. Never take it as your only treatment unless specifically directed.

3

CHECK ANTACIDS & MINERALS

Aluminum, magnesium, calcium, and iron can reduce absorption. Ask for exact timing.

4

CHECK RIFAMPIN & SEIZURE DRUGS

Some medicines can lower dolutegravir levels and may require a different schedule or alternative.

5

DOFETILIDE MUST NOT BE USED

This heart-rhythm medicine is contraindicated with Tivicay.

6

KEEP LABS & REFILLS ON TIME

Viral load, liver health, and the complete medication list must be reviewed regularly.

MY COMPLETE HIV REGIMEN

My Tivicay dose/time: _____

Other HIV medicine(s): _____

My HIV clinician: _____

MISSED A DOSE?

Take it when you remember unless the next dose is close. Do not double. Missing Tivicay or the companion medicines can weaken the whole regimen.

CALL RIGHT AWAY FOR

- Rash with fever, facial swelling, or trouble breathing
- Yellow skin or eyes, dark urine, or severe abdominal pain
- Severe weakness or persistent vomiting
- New depression or thoughts of self-harm
- Any gap in Tivicay or companion medicines

TIMING & INTERACTIONS

ANTACIDS

Aluminum or magnesium antacids generally require separation from Tivicay.

CALCIUM / IRON

May sometimes be taken together with food; otherwise separation is needed.

RIFAMPIN / SEIZURE DRUGS

These can change dolutegravir exposure. Never adjust the schedule yourself.

IMPORTANT SAFETY & SUCCESS REMINDERS

COMPLETE REGIMEN

Know every HIV medicine in your plan and refill them together.

LIVER HEALTH

Extra monitoring may be needed with hepatitis B or C.

PREGNANCY PLANNING

Tell the HIV clinician promptly about pregnancy or plans.

PRIVACY & SUPPORT

Fairview checks the full regimen discreetly and prevents refill gaps.



MY TIVICAY® GUIDE

KEEP THIS PAGE WITH YOUR MEDICATIONS

MY TREATMENT PLAN

My name: _____

My Tivicay time: _____

Other HIV medicines: _____

My HIV clinician: _____

Next refill date: _____

TIVICAY IS NOT A COMPLETE REGIMEN ALONE

IF I MISS A DOSE

- 1 Take it when remembered unless the next dose is close.
- 2 Do not take two doses together.
- 3 Check whether companion HIV medicines were also missed.
- 4 Call early if any part of the regimen may run out.

CHECK BEFORE TAKING

ANTACIDS

Magnesium or aluminum products need separation.

CALCIUM / IRON

Food changes the timing rule; confirm first.

RIFAMPIN

May require a different Tivicay schedule.

DOFETILIDE

Must not be used with Tivicay.

CALL MY CARE TEAM FOR

- Rash, fever, or facial swelling
- Yellow skin or eyes or dark urine
- Severe weakness or persistent vomiting
- New depression or self-harm thoughts
- A new prescription or supplement
- Any part of the regimen running out

KNOW MY WHOLE REGIMEN

Tivicay treats HIV only when used with the appropriate companion medicine or medicines. Missing one part can compromise the plan.

REFILL ALL REGIMEN COMPONENTS ON TIME

FAIRVIEW SUPPORT

PRIVATE COUNSELING

Questions answered discreetly.

INTERACTION CHECKS

Every prescription and supplement reviewed.

REFILL PROTECTION

Call early if travel, insurance, or supply may cause a gap.

COST SUPPORT

Coverage and assistance options checked for your prescription.